

Derivation of Continuous Linear Multistep Hybrid Block Method for the Solution of Volterra Integral Equation of Second Kind

Abstracts

In this paper, we proposed a continuous linear multistep hybrid block method for the solution of Volterra integral equation of second kind of the form $y(x) = f(x) + \int_{x_0}^{x_n} \varphi(x, s)y(s)ds$, using power series and trigonometrically fitted function as the trial solution for the approximation via collocation techniques. The proposed hybrid block scheme is found to be consistent, zero-stable and convergent. The implementation of the scheme on numerical problems and comparisons of results obtained with existing numerical method will be included

Keywords: Multistep hybrid block method, power series, collocation and interpolation method, second kind of Volterra integral equations.

1. Introduction

Volterra integral equation is a special kind of integral equation which is classified into three: the first, second and third kind. In this research, we develop two off - grid points of hybrid block method for the solution of second kind of Volterra integral equation due to its characteristics and uniqueness. In the literature, the second kind of Volterra integral equation (VIE) according to [1] is of the form: $y(x) = f(x) + \int_{x_0}^{x_n} \varphi(x, s)y(s)ds$ (1)

Where $f(x)$ is a given function and $\varphi(x, s)$ is called the kernel of integral equation. Volterra integral equation (VIE) appears especially when we try to transform an initial value problem into integral form, so that, the solution of the equation can be easily obtained than the original initial value problem [2]. Solving (1) is equivalent to solving the following initial value problem for ordinary differential equations of the first order

$$y'(x) = f'(x) + \varphi(x, y(x)), y(x_0) = f(x_0) \quad (2)$$

The Volterra Integral Equations are widely used in population growth models, physics, chemistry, and engineering [3]. Particularly important in science and engineering are systems of linear integral equations and their precise or approximate solutions. In the domains of engineering and applied research, some of these integral equations cannot be solved explicitly, hence approximation or numerical methods must frequently be used [4]. In recent years, many

strategies for resolving Volterra integral equations are suggested by many researchers such as: [5], [6], [7], [8], and much recently by [9], [10], [11], [12], [13],[14] and [15] respectively.

2. Derivation of the proposed method

In this section, we derive three step with two off-grid points of hybrid block method for the integration of Volterra integral equation of second kind by carefully selecting $p = \frac{1}{3}$ and $q = \frac{2}{3}$ for $p, q \in [0,1]$

Let the approximate series solution and trigonometrically fitted function of the Eq. (1) takes the

$$\text{form of } z(x) = \sum_{j=0}^3 \phi_j x^j + \sum_{j=1}^2 \lambda_j \sin x + \sum_{j=1}^2 \lambda_j \cos x \quad (3)$$

Where ϕ_j and λ_j are the coefficients to be determined.

Consider the ordinary differential equation

$$z'' = f(x, z, z'), z(a_0) = z_0, z'(a) = z'_0 \quad (4)$$

Subject to the condition

$$z(x) = y(x) - f(x) \quad (5)$$

The second derivative of Eq. (3) is given as;

$$z''(x) = \sum_{j=0}^3 j(j-1)\phi_j x^{j-2} - \sum_{j=1}^2 \lambda_j (\sin x) - \sum_{j=1}^2 \lambda_j (\cos x) \quad (6)$$

Substituting Eq. (4) into (1) gives

$$g(x, z, z') = \sum_{j=0}^3 j(j-1)\phi_j x^{j-2} - \sum_{j=1}^2 \lambda_j \sin x - \sum_{j=1}^2 \lambda_j \cos x \quad (7)$$

Interpolating (3) at $x_{n+\varpi}, \varpi = 0,1$ and collocating (6) at $x_{n+\vartheta}, \vartheta = \{0, \frac{1}{3}, \frac{2}{3}, 1, 2, 3\}$ leads to the

system of nonlinear equations written in the form

$$z_n(x) = X(x)A \quad (8)$$

$$\begin{bmatrix} 3 & 4x_n & -\frac{3}{2}x_n^2 & -\frac{1}{2}x_n^3 & \frac{17}{24}x_n^4 & \frac{11}{40}x_n^5 & -\frac{4}{45}x_n^6 & -\frac{8}{315}x_n^7 \\ 3(4x_n+4h) & -\frac{3}{2}(x_n+h)^2 & -\frac{1}{2}(x_n+h)^3 & \frac{17}{24}(x_n+h)^4 & \frac{11}{40}(x_n+h)^5 & -\frac{4}{45}(x_n+h)^6 & -\frac{8}{315}(x_n+h)^7 \\ 0 & 0 & -3 & -3x_n & \frac{17}{2}x_n^2 & \frac{11}{2}x_n^3 & -\frac{8}{3}x_n^4 & -\frac{16}{15}x_n^5 \\ 0 & 0 & -3 & -3x_n-h & \frac{17}{2}\left(x_n+\frac{1}{3}h\right)^2 & \frac{11}{2}\left(x_n+\frac{1}{3}h\right)^3 & -\frac{8}{3}\left(x_n+\frac{1}{3}h\right)^4 & -\frac{16}{15}\left(x_n+\frac{1}{3}h\right)^5 \\ 0 & 0 & -3 & -3x_n-2h & \frac{17}{2}\left(x_n+\frac{2}{3}h\right)^2 & \frac{11}{2}\left(x_n+\frac{2}{3}h\right)^3 & -\frac{8}{3}\left(x_n+\frac{2}{3}h\right)^4 & -\frac{16}{15}\left(x_n+\frac{2}{3}h\right)^5 \\ 0 & 0 & -3 & -3x_n-3h & \frac{17}{2}(x_n+h)^2 & \frac{11}{2}(x_n+h)^3 & -\frac{8}{3}(x_n+h)^4 & -\frac{16}{15}(x_n+h)^5 \\ 0 & 0 & -3 & -3x_n-6h & \frac{17}{2}(x_n+2h)^2 & \frac{11}{2}(x_n+2h)^3 & -\frac{8}{3}(x_n+2h)^4 & -\frac{16}{15}(x_n+2h)^5 \\ 0 & 0 & -3 & -3x_n-9h & \frac{17}{2}(x_n+3h)^2 & \frac{11}{2}(x_n+3h)^3 & -\frac{8}{3}(x_n+3h)^4 & -\frac{16}{15}(x_n+3h)^5 \end{bmatrix} \begin{bmatrix} \psi_0 \\ \psi_1 \\ \phi_0 \\ \phi_{\frac{1}{3}} \\ \phi_{\frac{2}{3}} \\ \phi_1 \\ \phi_2 \\ \phi_3 \end{bmatrix} = \begin{bmatrix} z_n \\ z_{n+1} \\ g_n \\ g_{n+\frac{1}{3}} \\ g_{n+\frac{2}{3}} \\ g_{n+1} \\ g_{n+2} \\ g_{n+3} \end{bmatrix}$$

Using the Gaussian elimination method to solve Eq. (8) gives the coefficients

$\psi_0, \psi_1, \phi_0, \phi_{\frac{1}{3}}, \phi_{\frac{2}{3}}, \phi_1, \phi_2, \phi_3$, which are then substituted into (3) and simplified to give the implicit

second derivative hybrid block method of the form;

$$z(x) = \sum_{i=0,1} \psi_i z_{n+i} + h^2 \left[\sum_{i=\frac{1}{3}, \frac{2}{3}} \phi_i g_{n+i} + \sum_{i=0}^3 \phi_i g_{n+i} \right] \quad (9)$$

Differentiating equation (9) to give:

$$z'(x) = \frac{1}{h} \sum_{i=0,1} \psi_i z_{n+i} + h \left[\sum_{i=\frac{1}{3}, \frac{2}{3}} \phi_i g_{n+i} + \sum_{i=0}^3 \phi_i g_{n+i} \right] \quad (10)$$

Where

$$\psi_0 = 1 - \frac{-x_n + x}{h} :$$

$$\psi_1 = \frac{-x_n + x}{h} :$$

$$\begin{aligned}
& -\frac{27}{40} \frac{(-x_n + x)^6}{h^4} + \frac{81}{1120} \frac{(-x_n + x)^7}{h^5} \phi_0 = -\frac{32}{315} (-x_n + x) h + \frac{1}{2} (-x_n + x)^2 - \frac{19}{18} \frac{(-x_n + x)^3}{h} \\
& + \frac{55}{48} \frac{(-x_n + x)^4}{h^2} - \frac{31}{48} \frac{(-x_n + x)^5}{h^3} \\
& + \frac{7}{40} \frac{(-x_n + x)^6}{h^4} - \frac{1}{56} \frac{(-x_n + x)^7}{h^5} \\
\phi_{\frac{1}{3}} = & -\frac{729}{2240} (-x_n + x) h + \frac{81}{40} \frac{(-x_n + x)^3}{h} - \frac{27}{8} \frac{(-x_n + x)^4}{h^2} + \frac{729}{320} \frac{(-x_n + x)^5}{h^3}
\end{aligned}$$

$$\begin{aligned}
\phi_{\frac{2}{3}} = & -\frac{81}{1960} (-x_n + x) h - \frac{81}{56} \frac{(-x_n + x)^3}{h} + \frac{783}{224} \frac{(-x_n + x)^4}{h^2} - \frac{3159}{1120} \frac{(-x_n + x)^5}{h^3} \\
& + \frac{513}{560} \frac{(-x_n + x)^6}{h^4} - \frac{81}{784} \frac{(-x_n + x)^7}{h^5}
\end{aligned}$$

$$\begin{aligned}
\phi_1 = & -\frac{11}{336} (-x_n + x) h + \frac{1}{2} \frac{(-x_n + x)^3}{h} - \frac{4}{3} \frac{(-x_n + x)^4}{h^2} + \frac{101}{80} \frac{(-x_n + x)^5}{h^3} - \frac{9}{20} \frac{(-x_n + x)^6}{h^4} \\
& + \frac{3}{56} \frac{(-x_n + x)^7}{h^5}
\end{aligned}$$

$$\begin{aligned}
\phi_2 = & \frac{1}{840} (-x_n + x) h - \frac{1}{40} \frac{(-x_n + x)^3}{h} + \frac{7}{96} \frac{(-x_n + x)^4}{h^2} - \frac{13}{160} \frac{(-x_n + x)^5}{h^3} + \frac{3}{80} \frac{(-x_n + x)^6}{h^4} \\
& - \frac{3}{560} \frac{(-x_n + x)^7}{h^5}
\end{aligned}$$

$$\begin{aligned}
\phi_3 = & -\frac{13}{141120} (-x_n + x) h + \frac{1}{504} \frac{(-x_n + x)^3}{h} - \frac{1}{168} \frac{(-x_n + x)^4}{h^2} + \frac{47}{6720} \frac{(-x_n + x)^5}{h^3} \\
& - \frac{1}{280} \frac{(-x_n + x)^6}{h^4} + \frac{1}{1568} \frac{(-x_n + x)^7}{h^5}
\end{aligned}$$

Evaluating Eq. (9) at non-interpolating points

$x = x_{n+\frac{1}{3}}, x_{n+\frac{2}{3}}, x_{n+2}, x_{n+3}$ yields

$$\begin{aligned}
z_{n+\frac{1}{3}} = & \frac{2}{3} z_n + \frac{1}{3} z_{n+1} - \frac{497}{87480} h^2 g_n - \frac{1}{15} h^2 g_{n+\frac{1}{3}} - \frac{523}{15120} h^2 g_{n+\frac{2}{3}} \\
& - \frac{31}{7290} h^2 g_{n+1} + \frac{1}{11664} h^2 g_{n+2} - \frac{1}{153090} h^2 g_{n+3}
\end{aligned}$$

$$\begin{aligned}
z_{n+\frac{2}{3}} = & \frac{1}{3} z_n + \frac{2}{3} z_{n+1} - \frac{29}{10935} h^2 g_n - \frac{167}{4320} h^2 g_{n+\frac{1}{3}} - \frac{79}{1260} h^2 g_{n+\frac{2}{3}} \\
& - \frac{209}{29160} h^2 g_{n+1} + \frac{1}{14580} h^2 g_{n+2} - \frac{11}{2449440} h^2 g_{n+3}
\end{aligned} \tag{11}$$

$$z_{n+3} = -2z_n + 3z_{n+1} + \frac{11}{120}h^2g_n + \frac{243}{560}h^2g_{n+\frac{2}{3}} + \frac{13}{10}h^2g_{n+1} + \frac{89}{80}h^2g_{n+2} + \frac{13}{210}h^2g_{n+3}$$

$$h z_n' = -z_n + z_{n+1} - \frac{32}{315} h^2 g_n - \frac{729}{2240} h^2 g_{n+\frac{1}{3}} - \frac{81}{1960} h^2 g_{n+\frac{2}{3}} - \frac{11}{336} h^2 g_{n+1} \\ + \frac{1}{840} h^2 g_{n+2} - \frac{13}{141120} h^2 g_{n+3}$$

$$hz'_{n+\frac{2}{3}} = -z_n + z_{n+1} + \frac{131}{25515} h^2 g_n + \frac{547}{4032} h^2 g_{n+\frac{1}{3}} + \frac{839}{17640} h^2 g_{n+\frac{2}{3}} - \frac{2999}{136080} h^2 g_{n+1} + \frac{5}{13608} h^2 g_{n+2} - \frac{269}{11430720} h^2 g_{n+3}$$

$$h z'_{n+2} = -z_n + z_{n+1} - \frac{53}{315} h^2 g_n + \frac{459}{448} h^2 g_{n+\frac{1}{3}} - \frac{621}{392} h^2 g_{n+\frac{2}{3}} + \frac{3193}{1680} h^2 g_{n+1} \\ + \frac{281}{840} h^2 g_{n+2} - \frac{1021}{141120} h^2 g_{n+3}$$

(12)

This gives the following equation in matrix form

$$A_1 Z = A_2 R_1 + B_1 R_2 + B_2 R_3 \quad (13)$$

$$A_1 = \begin{bmatrix} 1 & 0 & -\frac{1}{3} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -\frac{2}{3} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -2 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -3 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -\frac{1}{h} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -\frac{1}{h} & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & -\frac{1}{h} & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & -\frac{1}{h} & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & -\frac{1}{h} & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, z = \begin{bmatrix} z_{n+\frac{1}{3}} \\ z_{n+\frac{2}{3}} \\ z_{n+1} \\ z_{n+2} \\ z_{n+3} \\ z'_{n+\frac{1}{3}} \\ z'_{n+\frac{2}{3}} \\ z'_{n+1} \\ z'_{n+2} \\ z'_{n+3} \end{bmatrix}, A_2 = \begin{bmatrix} \frac{2}{3} & 0 \\ -\frac{1}{3} & 0 \\ \frac{1}{3} & 0 \\ 1 & 0 \\ 2 & 0 \\ 1 & 0 \\ h & 1 \\ \frac{1}{h} & 0 \\ \frac{1}{h} & 0 \\ \frac{1}{h} & 0 \end{bmatrix}, R_1 = \begin{bmatrix} z_n \\ z'_n \end{bmatrix}, B_1 = \begin{bmatrix} -\frac{497}{h^2} \\ 87480 \\ \frac{29}{h^2} \\ 10935 \\ \frac{1}{h^2} \\ 15 \\ \frac{11}{h^2} \\ 120 \\ \frac{32}{h} \\ 315 \\ 5687 \\ 408240 \\ \frac{131}{h} \\ 15515 \\ \frac{11}{h} \\ 1008 \\ \frac{53}{h} \\ 315 \\ 3079 \\ 5040 \end{bmatrix}, R_2 = [g_n], B_2 = \begin{bmatrix} -\frac{1}{h^2} & -\frac{53}{h^2} & -\frac{31}{h^2} & \frac{1}{h^2} & -\frac{1}{h^2} \\ 15 & 15120 & 7290 & 11664 & 153090 \\ -\frac{167}{h^2} & -\frac{79}{h^2} & -\frac{209}{h^2} & \frac{1}{h^2} & -\frac{11}{h^2} \\ 4320 & 1260 & 29160 & 14580 & 2449440 \\ \frac{81}{h^2} & -\frac{81}{h^2} & \frac{127}{h^2} & \frac{1}{h^2} & -\frac{3}{h^2} \\ 160 & 140 & 120 & 12 & 1120 \\ 0h^2 & \frac{243}{h^2} & \frac{13}{h^2} & \frac{89}{h^2} & \frac{13}{h^2} \\ & 560 & 10 & 80 & 210 \\ & -\frac{729}{h^2} & -\frac{81}{h^2} & -\frac{11}{h^2} & -\frac{1}{h^2} \\ 2240 & 1960 & 336 & 840 & 141120 \\ -\frac{13}{h^2} & -\frac{2231}{h^2} & \frac{127}{h^2} & -\frac{53}{h^2} & -\frac{131}{h^2} \\ 504 & 14112 & 34020 & 1428840 & 272160 \\ -\frac{547}{h^2} & -\frac{839}{h^2} & -\frac{2999}{h^2} & \frac{5}{h^2} & -\frac{269}{h^2} \\ 4032 & 17640 & 136080 & 13608 & 14430720 \\ \frac{27}{h^2} & \frac{2133}{h^2} & \frac{17}{h^2} & -\frac{1}{h^2} & \frac{1}{h^2} \\ 280 & 7840 & 140 & 1120 & 17640 \\ \frac{459}{h^2} & -\frac{621}{h^2} & \frac{3193}{h^2} & \frac{281}{h^2} & -\frac{1021}{h^2} \\ 448 & 392 & 1680 & 840 & 141120 \\ -\frac{729}{h^2} & \frac{35397}{h^2} & -\frac{817}{h^2} & \frac{1097}{h^2} & \frac{1025}{h^2} \\ 280 & 7840 & 420 & 672 & 3528 \end{bmatrix}, R_3 = \begin{bmatrix} g_1 \\ g_2 \\ g_3 \end{bmatrix} \quad (14)$$

Substituting Eq.(14) into Eq. (13) and multiply by the inverse of A_1 gives the hybrid block in the form:

$$IZ_1 = \bar{B} R_1 + \bar{D}_1 R_2 + \bar{D}_2 R_3 \quad (15)$$

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} z_{m+3}^1 \\ z_{m+3}^2 \\ z_{m+1} \\ z_{m+2} \\ z_{m+3} \\ z_{m+3}^1 \\ z_{m+3}^2 \\ z_{m+1} \\ z_{m+2} \\ z_{m+3} \end{bmatrix} = \begin{bmatrix} 1 & \frac{1}{3}h \\ 1 & \frac{2}{3}h \\ 1 & h \\ 1 & 2h \\ 1 & 3h \\ 0 & 1 \\ 0 & 1 \\ 0 & 1 \\ 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} z_n \\ z_n \end{bmatrix} + \begin{bmatrix} \frac{17257}{h^2} \\ 612360 \\ \frac{4981}{h^2} \\ 76545 \\ \frac{32}{h^2} \\ 315 \\ \frac{43}{h^2} \\ 315 \\ \frac{11}{h^2} \\ 280 \\ \frac{6737}{h} \\ 58320 \\ \frac{389}{h} \\ 3645 \\ 9 \\ -h \\ 80 \\ \frac{1}{h} \\ 15 \\ \frac{57}{h} \\ 80 \end{bmatrix} + \begin{bmatrix} \frac{281}{h^2} & \frac{2203}{h^2} & \frac{2719}{h^2} & \frac{127}{h^2} & \frac{829}{h^2} \\ 6780 & 105840 & 408240 & 408240 & 34292160 \\ \frac{337}{h^2} & \frac{31}{h^2} & \frac{374}{h^2} & \frac{37}{h^2} & \frac{61}{h^2} \\ 1890 & 882 & 25515 & 51030 & 1071630 \\ \frac{729}{h^2} & \frac{281}{h^2} & \frac{11}{h^2} & \frac{1}{h^2} & \frac{13}{h^2} \\ 2240 & 6780 & 336 & 840 & 141120 \\ \frac{81}{h^2} & \frac{243}{h^2} & \frac{118}{h^2} & \frac{17}{h^2} & \frac{11}{h^2} \\ 70 & 490 & 105 & 210 & 4410 \\ \frac{2187}{h^2} & \frac{2187}{h^2} & \frac{783}{h^2} & \frac{621}{h^2} & \frac{195}{h^2} \\ 2240 & 3190 & 560 & 560 & 3136 \\ \frac{826}{h} & \frac{1177}{h} & \frac{709}{h} & \frac{206}{h} & \frac{1483}{h} \\ 2880 & 10080 & 19440 & 178605 & 3810240 \\ 83 & 4 & 13 & 1 & 1 \\ -h & -h & -h & -h & -h \\ 180 & 45 & 1215 & 1215 & 14580 \\ 27 & 351 & 37 & 1 & 1 \\ -h & -h & -h & -h & -h \\ 64 & 1120 & 240 & 480 & 6720 \\ 27 & 54 & 29 & 1 & 1 \\ -h & -h & -h & -h & -h \\ 20 & 35 & 15 & 3 & 140 \\ \frac{729}{h} & \frac{729}{h} & \frac{153}{h} & \frac{261}{h} & \frac{93}{h} \\ 320 & 160 & 80 & 160 & 320 \end{bmatrix} \begin{bmatrix} g_1 \\ g_2 \\ g_3 \\ g_1 \\ g_2 \\ g_3 \end{bmatrix}$$

By putting Eq. (5) in Eq. (15) yield the proposed hybrid blockschemes of the form:

$$\begin{aligned}
y_{n+\frac{1}{3}} - f_{n+\frac{1}{3}} &= y_n - f_n + \frac{1}{3} h(y'_n - f'_n) + \frac{17257}{612360} h^2 g_n + \frac{281}{6720} h^2 g_{n+\frac{1}{3}} - \frac{2203}{105840} h^2 g_{n+\frac{2}{3}} \\
&+ \frac{2719}{408240} h^2 g_{n+1} - \frac{127}{408240} h^2 g_{n+2} + \frac{829}{34292160} h^2 g_{n+3} \\
y_{n+\frac{2}{3}} - f_{n+\frac{2}{3}} &= y_n - f_n + \frac{2}{3} h(y'_n - f'_n) + \frac{4981}{76545} h^2 g_n + \frac{337}{1890} h^2 g_{n+\frac{1}{3}} - \frac{31}{882} h^2 g_{n+\frac{2}{3}} + \\
&\frac{374}{25515} h^2 g_{n+1} - \frac{37}{51030} h^2 g_{n+2} + \frac{61}{1071630} h^2 g_{n+3} \\
y_{n+1} - f_{n+1} &= y_n - f_n + h(y'_n - f'_n) + \frac{32}{315} h^2 g_n + \frac{729}{2240} h^2 g_{n+\frac{1}{3}} + \frac{81}{1960} h^2 g_{n+\frac{2}{3}} + \frac{11}{336} h^2 g_{n+1} \\
&- \frac{1}{840} h^2 g_{n+2} + \frac{13}{141120} h^2 g_{n+3} \\
y_{n+2} - f_{n+2} &= y_n - f_n + 2h(y'_n - f'_n) + \frac{43}{315} h^2 g_n + \frac{81}{70} h^2 g_{n+\frac{1}{3}} - \frac{243}{490} h^2 g_{n+\frac{2}{3}} + \frac{118}{105} h^2 g_{n+1} \\
&+ \frac{17}{210} h^2 g_{n+2} - \frac{11}{4410} h^2 g_{n+3} \\
y_{n+3} - f_{n+3} &= y_n - f_n + 3h(y'_n - f'_n) + \frac{111}{280} h^2 g_n + \frac{2187}{2240} h^2 g_{n+\frac{1}{3}} + \frac{2187}{3920} h^2 g_{n+\frac{2}{3}} + \frac{783}{560} h^2 g_{n+1} \\
&+ \frac{621}{560} h^2 g_{n+2} + \frac{195}{3136} h^2 g_{n+3} \\
y'_{n+\frac{1}{3}} - f'_{n+\frac{1}{3}} &= y'_n - f'_n + \frac{6737}{58320} h g_n + \frac{863}{2880} h g_{n+\frac{1}{3}} - \frac{1177}{10080} h g_{n+\frac{2}{3}} + \frac{709}{19440} h g_{n+1} \\
&- \frac{206}{178605} h g_{n+2} - \frac{1483}{3810240} h g_{n+3} \\
y'_{n+\frac{2}{3}} - f'_{n+\frac{2}{3}} &= y'_n - f'_n + \frac{389}{3645} h g_n + \frac{83}{180} h g_{n+\frac{1}{3}} + \frac{4}{45} h g_{n+\frac{2}{3}} + \frac{13}{1215} h g_{n+1} \\
&- \frac{1}{1215} h g_{n+2} + \frac{1}{14580} h g_{n+3} \\
y'_{n+1} - f'_{n+1} &= y'_n - f'_n + \frac{9}{80} h g_n + \frac{27}{64} h g_{n+\frac{1}{3}} + \frac{351}{1120} h g_{n+\frac{2}{3}} + \frac{37}{240} h g_{n+1}
\end{aligned} \tag{16}$$

$$- \frac{1}{480} h g_{n+2} + \frac{1}{6720} h g_{n+3}$$

$$y'_{n+2} - f'_{n+2} = y'_n - f'_n - \frac{1}{15} h g_n + \frac{27}{20} h g_{n+\frac{1}{3}} - \frac{54}{35} h g_{n+\frac{2}{3}} + \frac{29}{15} h g_{n+1} + \frac{1}{3} h g_{n+2}$$

$$- \frac{1}{140} h g_{n+3}$$

$$y'_{n+3} - f'_{n+3} = y'_n - f'_n + \frac{57}{80} h g_n - \frac{729}{320} h g_{n+\frac{1}{3}} + \frac{729}{160} h g_{n+\frac{2}{3}} - \frac{153}{80} h g_{n+1} + \frac{261}{160} h g_{n+2}$$

$$+ \frac{93}{320} h g_{n+3}$$

3. Analysis of the Hybrid Block Method

In this section, the analysis of the order, error constant, convergence and stability of proposed scheme is carried out.

3.1. Order and error constant of the proposed method

Let the linear difference operator ℓ associated with the new method (16) be defined as

$$L[y(x; h)] = \sum_{j=0}^k \alpha_j y(x + jh) - h^2 \sum_{j=0}^k (\beta_j y''(x + jh)) \quad (17)$$

Where $y(x)$ is an arbitrary test function continuously differential on $[a, b]$ Expanding $y(x + jh)$, $y'(x + jh)$ and $y''(x + jh)$ of (16) in Taylor series in the form:

$$\begin{aligned}
& \sum_{j=0}^{\infty} \frac{\left(\frac{1}{3}h\right)^j}{j!} (z_n)^j - z_n - \frac{1}{3}hz'_n - \sum_{j=0}^{\infty} \frac{h^{j+2}}{j!} g_n^{j+2} \left[\frac{1725}{612360} + \frac{281}{6720} \left(\frac{1}{3}\right) - \frac{2203}{1058040} \left(\frac{2}{3}\right) + \frac{2719}{408240} (1) - \frac{127}{408240} (2) + \frac{829}{3429460} (3) \right] \\
& \sum_{j=0}^{\infty} \frac{\left(\frac{2}{3}h\right)^j}{j!} (z_n)^j - z_n - \frac{2}{3}hz'_n - \sum_{j=0}^{\infty} \frac{h^{j+2}}{j!} g_n^{j+2} \left[\frac{4981}{76545} + \frac{337}{1890} \left(\frac{1}{3}\right) - \frac{31}{882} \left(\frac{2}{3}\right) + \frac{374}{25515} (1) - \frac{37}{51030} (2) + \frac{61}{1071630} (3) \right] \\
& \sum_{j=0}^{\infty} \frac{(h)^j}{j!} (z_n)^j - z_n - hz'_n - \sum_{j=0}^{\infty} \frac{h^{j+2}}{j!} g_n^{j+2} \left[\frac{32}{315} + \frac{729}{2240} \left(\frac{1}{3}\right) + \frac{81}{1960} \left(\frac{2}{3}\right) + \frac{11}{336} (1) - \frac{1}{840} (2) + \frac{13}{141120} (3) \right] \\
& \sum_{j=0}^{\infty} \frac{(2h)^j}{j!} (z_n)^j - z_n - 2hz'_n - \sum_{j=0}^{\infty} \frac{h^{j+2}}{j!} g_n^{j+2} \left[\frac{43}{315} + \frac{81}{70} \left(\frac{1}{3}\right) - \frac{243}{490} \left(\frac{2}{3}\right) + \frac{118}{105} (1) + \frac{17}{210} (2) - \frac{11}{4410} (3) \right] \\
& \sum_{j=0}^{\infty} \frac{(3h)^j}{j!} (z_n)^j - z_n - 3hz'_n - \sum_{j=0}^{\infty} \frac{h^{j+2}}{j!} g_n^{j+2} \left[\frac{111}{280} + \frac{2187}{2240} \left(\frac{1}{3}\right) + \frac{2187}{3920} \left(\frac{2}{3}\right) + \frac{783}{560} (1) + \frac{621}{660} (2) + \frac{195}{3136} (3) \right] \\
& \sum_{j=0}^{\infty} \frac{\left(\frac{1}{3}\right)^j}{j!} (z_n)^j - z_n - \sum_{j=0}^{\infty} \frac{h^{j+1}}{j!} g_n^{j+2} \left[\frac{6737}{58320} + \frac{863}{2880} \left(\frac{1}{3}\right) - \frac{1177}{10080} \left(\frac{2}{3}\right) + \frac{709}{19440} (1) - \frac{13}{7776} (2) + \frac{211}{1632960} (3) \right] \\
& \sum_{j=0}^{\infty} \frac{\left(\frac{2}{3}\right)^j}{j!} (z_n)^j - z_n - \sum_{j=0}^{\infty} \frac{h^{j+1}}{j!} g_n^{j+2} \left[\frac{389}{3645} + \frac{83}{180} \left(\frac{1}{3}\right) + \frac{4}{45} \left(\frac{2}{3}\right) + \frac{13}{125} (1) - \frac{1}{1215} (2) + \frac{1}{14580} (3) \right] \\
& \sum_{j=0}^{\infty} \frac{(1)^j}{j!} (z_n)^j - z_n - \sum_{j=0}^{\infty} \frac{h^{j+1}}{j!} g_n^{j+2} \left[\frac{9}{80} + \frac{27}{64} \left(\frac{1}{3}\right) + \frac{351}{1120} \left(\frac{2}{3}\right) + \frac{37}{240} (1) - \frac{1}{480} (2) + \frac{1}{6720} (3) \right] \\
& \sum_{j=0}^{\infty} \frac{(2)^j}{j!} (z_n)^j - z_n - \sum_{j=0}^{\infty} \frac{h^{j+1}}{j!} g_n^{j+2} \left[\frac{-1}{15} + \frac{27}{20} \left(\frac{1}{3}\right) - \frac{54}{35} \left(\frac{2}{3}\right) + \frac{29}{15} (1) + \frac{1}{3} (2) - \frac{1}{140} (3) \right] \\
& \sum_{j=0}^{\infty} \frac{(3)^j}{j!} (z_n)^j - z_n - \sum_{j=0}^{\infty} \frac{h^{j+1}}{j!} g_n^{j+2} \left[\frac{57}{80} - \frac{729}{160} \left(\frac{1}{3}\right) + \frac{729}{160} \left(\frac{2}{3}\right) - \frac{153}{80} (1) + \frac{261}{160} (2) + \frac{93}{320} (3) \right]
\end{aligned}
\tag{18}$$

If we assume that $y(x)$ has many higher derivatives and collecting the terms, we have:

$$\ell[y(x); h] = \bar{c}_0 y(x) + \bar{c}_1 h y'(x) + \bar{c}_2 h^2 y''(x) + \dots + \bar{c}_{p+2} h^{p+2} y^{(p+2)}(x) \tag{19}$$

According to [11], the proposed scheme has order p if $\bar{c}_0 = \bar{c}_1 = \bar{c}_2 = \dots = \bar{c}_{p+1} = 0, \bar{c}_{p+2} \neq 0$

The proposed method is of order $p = [7, 7, 7, 7, 7, 7, 7, 7, 7]^T$, with error constant

C_9

$$= \left[\frac{286}{793618560}, \frac{107}{12400290}, \frac{1}{72576}, \frac{1}{5670}, \frac{11}{13440}, \frac{281}{14696640}, \frac{281}{2755620}, \frac{11}{544320}, \frac{17}{34020}, \frac{13}{4032} \right]^T$$

3.2. Consistency of the method

According to Areo and Omojola (2015), the hybrid block method is said to be consistent if it has an order more than or equal to one, i.e. ($p \geq 1$). Since the Eq. (16) is of order $p = 7$, therefore, the proposed hybrid block method is consistent

2.3. Zero-stability of the proposed method

The linear multistep hybrid block method is said to be zero-stable as $h \rightarrow 0$, if the roots of the first characteristics polynomial defined by $\rho(z) = \det[\sum_{j=0}^k A^{(i)} z^{(K-i)}]$ satisfies $|z| \leq 1$ and every root of $|z| = 1$ has multiplicity not exceeding the order of the differential equation. Awoyemi *et.al.* (2011)

$$A^0 = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \quad A' = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

We have $\rho(z) = z^8(z - 1) = 0, z = 0.0.0.0.0.0.0, 1$

Since $|z| = 1$, therefore. The proposed method is zero-stable

2.4. Convergence of the proposed method The necessary and sufficient condition for a linear multistep hybrid block method to be convergent is to be consistent and zero stable. Since, our scheme satisfies the two conditions, hence the Eq. (9) is convergent.

4. Implementation of method

In this section, we implement the proposed hybrid block method on two considered problems of second kind VIEs.

Problem 4.1

Consider the second kind linear volterra integral equation

$$X(t) = t^2 + \int_0^t (t-s)x_1(s)ds$$

With exact solution $X(t) = 2\cos ht - 2, h = 0.1$

Source: Muturiet al. (2014)

Converting the nonlinear VIE to a second order ODE, we obtained

$$t^2 + \int_0^t (t-s)x_1(s)ds$$

$$\int_0^t x_1(s) ds$$

$$x(t) =$$

$$x'(t) = 2t +$$

$$x''(t) = 2 + x_1(s)$$

Then, the second order ODE is given as

$$z'' = 0z_0' + z_0 + 2$$

Where $z_0 = 0, z' = 0$.

Table .1 the exact solution and computed results from the propose methods for problem 1

X	Exact	Numerical Result	Error in Muturiet	Error in Proposed
---	-------	------------------	-------------------	-------------------

			al. (2014)	Method
0.1	0.01000833611160719800	0.01001667222375875757	1.0 E-05	2.7763 E-13
0.2	0.04013351123815169260	0.04013351123459477332	3.0 E-05	3.5569 E-12
0.3	0.09067702825772097000	0.09067702827425125113	8.0 E-05	1.6530 E-11
0.4	0.16214474367690961860	0.16214474375907952666	1.40 E-04	8.2169 E-11
0.5	0.25525193041276157040	0.25525193055694029498	2.20 E-04	1.4417 E-10
0.6	0.37093043648453540760	0.37093043671821427031	3.20 E-04	2.33678 E-10
0.7	0.51033801126188603640	0.51033801163676863682	4.40 E-04	3.74882 E-10
0.8	0.67486989260968919600	0.67486989312433225989	5.90E-04	5.14643 E-10
0.9	0.86617277089754877560	0.86617277158760432182	7.70E-04	6.9005 E-10
1.0	1.08616126963048755700	1.08616127056068089180	9.80 E-04	9.3019 E-10

Problem 2
second kind linear volterra integral equation

$$U(x) = 1 + x + \int_0^x (x - t)U(t)dt$$

exact solution:

$$U(x) = e^x, \quad h = 0.1$$

Source:Shoukralla and Ahmed (2020)

VIE to a second order ODE, we obtained

$$\int_0^x (x - t) u(t)dt$$

$$u'(x) = 1 + \int_0^x u(t) dt u''(x) = u(x), \text{ The second order ODE is then given as } z'' = 0z' + z$$

$$\text{where } z_0 = 0, z' = 0.$$

exact solution and computed results from the propose methods for problem 2

Consider the

With

Converting the

$$u(x)1 + x +$$

Table 2 the

X	Exact	Numerical Result	Error in Shoukralla and Ahmed (2020)	Error in Proposed Method
0.1	1.1051709180756476248	1.10517091807580048840	1.4089 E-09	1.5286 E-13
0.2	1.2214027581601698339	1.22140275815817059020	9.1493 E-08	1.9992 E-12
0.3	1.3498588075760031040	1.34985880758546021050	1.0576 E-05	9.4571 E-12
0.4	1.4918246976412703178	1.49182469768812457810	6.0309 E-06	4.6854 E-11
0.5	1.6487212707001281468	1.64872127078173504370	2.3354 E-05	8.1607 E-11
0.6	1.8221188003905089749	1.82211880052608176310	7.08004 E-05	1.3557 E-10
0.7	2.0137527074704765216	2.01375270769626071760	1.8129E-04	2.2578 E-10
0.8	2.2255409284924676046	2.22554092880652021370	4.1026 E-04	3.1405 E-10
0.9	2.4596031111569496638	2.45960311158724651560	8.4486 E-04	4.3029 E-10
1.0	2.7182818284590452354	2.69286592330700708760	1.6151 E-03	2.5416 E-02

Discussion and Conclusion

In this research, Continuous Linear Multistep Hybrid Block Method was proposed for the solution of integral equation of second kind. We investigated the property of this proposed method in terms of order, error constant, consistency, zero-stability and convergence analysis. The scheme was also used to solve numerically two problems of volterra integral equation of second kind and the results were compared with [9] and [10]. From the Table 1 and 2, we discovered that, the proposed multistep hybrid block method were capable of Handling the second kind VIEs. The results obtained from Table 1 and 2 indicated that our proposed methods are considerably much more accurate than the existing numerical methods. All computation and program were carried out with the aid of MAPPLE 15 software.

References

1. Volterra, V. (1860). 1940. Obituary Notices of fellows of the royal society, 3(1941), 690-729
2. Gonzalez-Rodelas, P., Pasadas, M., Kouibia, A and Mustafa, B. (2022). Numerical solution of linear volterra integral equation system of second kind by Radial Basis function. *Mathematics*, 10(2), 223.
3. Wazwaz, A.M. Linear and Nonlinear Integral Equations: Methods and Applications, Springer **2011**.
4. Mechee, M.S., Al- Ramahi, A.M and Kadum, R.M. (2016). Variation iteration method for solving a class of volterra integral equations. *Journals of University of Babylon*, 24(9).
5. Baker, C.T.H. (2000) A perspective on the numerical treatment of Volterra equations, *Journal of Computer and Applied Mathematics*. 125 (1) 217–249.
6. Tahmasbi, A., and Fard, O.S. (2008). Numerical solution of linear Volterra integral equations system of the second kind, *Appl. Math. Comput.* 201 (1) 547–552.
7. Bhattacharya, S., & Mandal, B. N. (2008). Use of Bernstein polynomials is Numerical Solution of Volterra Integral Equations. *Applied Mathematical Sciences*, 2(36), 1773-1787
8. Ibrahimov, P. and Imanova, M. (2021). Multistep methods of the hybrid type and their application to solve the second kind volterra integral equation. *Symmetry*. 13, 1087-<https://doi.org/10.3390/sym13061087>.
9. Maturi, D.A., Bajamal, A.Z., and Algethami, B. (2014). Numerical solution of volterra IntegralEquation of the second kind using implicit trapezoidal. *Journal: Journal of advances in mathematics*, 8(2).
10. Shoukralla, E.S., and Ahmed, B.M. (2020). Numerical solutions of volterra integral equations of the second using Lagrange interpolation via thevandermonde matrix. In *Journal of Physics: conference series* (vol.1447, No.1, p.o12003).IOP published.
11. Aruchuman and Sulaiman, (2010) Non-standard Finite Difference Method for Numerical Solution of Second Order Linear FredholmIntegro-Differential Equations. <https://www.researchgate.net/publication/331716116>
12. Yahaya, Y. A. And Asabe A. T. (2016) A Block Hybrid Implicit Algorithms for Solution of First Order Differential Equations. *British Journal of Mathematics & Computer Science* 13(3): 1-9, 2016, Article no.BJMCS.19682 ISSN: 2231-0851.

13. Adu, A., Ine, Nyiutya Cephas and John., ChusehAhmadu (2023) Three-Step Three off Grid Points for Solving Volterra Integral Equation of the Second Kind. *Asian Journal of Pure and Applied Mathematics*, Vol. 5, Issue 1, Page 512-523; Article no. AJPAM.1409.
14. J.A. Chuseh, B.B. Rakiatou, A. Adu, A. Ibrahim and S.A. Kambai (2024), Application of Collocation Methods for a Hybrid Block Scheme for the Solution of Volterra Integral Equation of the Second Kind. *Asian Journal of Pure and Applied Mathematics*, Vol. 6, Issue 1, Page 118-131; Article No. AJPAM.1499.
15. Ahmed A. Khidir (2021) A New Numerical Technique for Solving Volterra Integral Equations Using Chebyshev Spectral Method.hindawi Mathematical problems in Engineering. Vol.2021, Article ID 9230714. <https://doi.org/10.1155/2021/9230714>
16. D. Raymond, A. Adu, P. O. Olanrewaju and R. Ajia (2023) Three-step Method of Volterra Integral Equation of the Second Kind.*Asian Journal of Pure and Applied Mathematics Volume 5, Issue 1, Page 264-273, 2023;Article no.AJPAM.1300*